

Giant thymolipoma in a child: The silent chest mass

S Otido,¹ MD; Z Dangor,¹ PhD; A Zanini,² MD; D Harrison,² MD

¹ Department of Paediatrics, Chris Hani Baragwanath Academic Hospital and Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

² Department of Paediatric Surgery, Chris Hani Baragwanath Academic Hospital and Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

Corresponding author: S Otido (samotido@gmail.com)

Thymolipoma is a rare benign anterior mediastinal tumour of thymic origin containing both thymic and mature adipose tissues. In most cases it has a silent course and can grow to large sizes before presenting with respiratory symptoms. We report a case of a giant thymolipoma in a 4-year-old girl treated at Chris Hani Baragwanath Academic Hospital, South Africa.

Afr J Thoracic Crit Care Med 2022;28(3):129-132. <https://doi.org/10.7196/AJTCCM.2022.v28i3.181>

Thymolipomas are rare benign anterior mediastinal tumours consisting of thymic and fatty tissues, and account for 2 - 9% of all thymic tumours.^[1] These tumours are slow growing and are usually diagnosed incidentally in older children and young adults. Surgical excision is the mainstay of treatment.^[1] We report the case of a young child with a massive thymolipoma who underwent successful surgical removal.

Case report

A 4-year-old HIV-uninfected girl was referred to the paediatric pulmonology clinic at Chris Hani Baragwanath Academic Hospital, South Africa (SA), with a 2-year history of chronic cough and poor weight gain. She had been treated with repeated courses of antibiotics for respiratory tract infections, and had also been started on tuberculosis (TB) therapy based on the chronic cough and abnormal findings on chest radiographs.

At presentation, she was stunted, had digital clubbing, and had dullness to percussion on the right side of the chest with reduced breath sounds. Her white cell count was raised ($20.7 \times 10^9/L$) with predominant lymphocytes (80.9%). The frontal chest radiograph showed a dense homogeneous opacification in the right middle and lower zones, and the left middle zone (Fig. 1). The lateral chest radiograph showed that the opacification was from the anterior mediastinum. Bronchoscopy showed a patent right upper lobe bronchus, but the right bronchus intermedius was completely occluded as a result of external compression.

The left upper and lower lobes were normal in structure with patent bronchi. A chest computed tomography (CT) scan (Fig. 2) showed an anterior soft-tissue hypodense mass on the right with herniation across the midline to the left hemithorax that displaced surrounding structures but was not encasing them. Histological examination of a Tru-Cut biopsy specimen showed benign thymic parenchyma with no evidence of neoplastic infiltration.

Sternotomy was performed and the entire mass was resected. The tumour was initially removed as a well-encapsulated bilobar mass, but the mass extended posteriorly into the right thorax and required some piecemeal resection (Fig. 3). Two large specimens were analysed, weighing 610 g and 360 g and measuring $170 \times 100 \times 30$ mm and $120 \times 60 \times 20$ mm, respectively. Microscopy revealed thymic parenchymal tissue mixed with mature adipose tissue, and no features of cytological atypia. Numerous Hassall's corpuscles were also seen. The final histopathology report confirmed that the tumour was a thymolipoma. There were no intra- or postoperative complications, and the patient was discharged.

Discussion

To our knowledge, our patient is the youngest child to have undergone surgical resection of a large thymolipoma in SA. Thymolipomas are rare, slow-growing benign mediastinal tumours that were first described by Hall in 1949.^[2] The global incidence has been reported as 0.12 per 100 000 cases per year.^[3] Thymolipoma has been linked to various autoimmune diseases,

with myasthenia gravis reported in half of the cases, but the pathogenesis remains unclear.

Other conditions that are associated with thymolipoma include red cell aplasia, Graves' disease, lichen planus and hypogammaglobulinaemia.^[3] The differential diagnoses for a large mediastinal mass include teratoma, thymoma, lipoma, lipomatosis, liposarcoma and thymoliposarcoma. Thymolipomas can occur at any age, although they are mainly found in young adults (mean age 33 years).^[4] However, to our knowledge only 36 cases have been reported in the current available English literature. The youngest patient ever reported was a 6-month-old infant who presented with respiratory distress due to compression of adjacent structures.^[5]

The chest radiograph can mimic abnormalities of adjacent structures, such as pericardial effusion, cardiomegaly, atelectasis or lung sequestration.^[6] Our patient was initially started on treatment for TB because of the nonspecific radiographic findings. Computed tomography and magnetic resonance imaging are usually diagnostic. They reveal a fatty mass with fibrous septa, which is highly suggestive of a thymolipoma.^[7] However, other differential diagnoses are possible, and they include potentially malignant diseases. Definitive diagnosis is through histopathological examination, which should demonstrate mature large lobules of adipose tissue mixed with normal thymus containing epithelial cells, thymic lymphocytes and Hassall's corpuscles.^[1] According to our literature

Table 1. Summary of a literature review of paediatric thymolipomas (references available as a supplementary file at <https://www.samedical.org/file/1873>)

Author	Age (yrs)	Presentation	Investigation	Biopsy	Treatment	Approach	Invasion	Malignancy	Complications
Maekawa <i>et al.</i> ^[51]	16	Incidental CXR, right bundle-branch block	CT chest + MRI	FNA CT guided	Total thymectomy	Sternotomy	No	No	No
Miranda <i>et al.</i> ^[52]	4	Recurrent LRTI, CXR	CT chest + MRI	No	Primary resection	n/d	No	No	No
Sundstrom ^[53]	8	Incidental CXR	No others	No	Primary resection, partial thymectomy	Thoracotomy	No	No	No
Damadoglu <i>et al.</i> ^[54]	16	Fatigue + myasthenia gravis	CT scan	No	Thymectomy	Sternotomy	No	No	No
Daneshbod ^[55]	17	Dyspnoea on exertion	CXR + CT	No	Thymectomy	Thoracotomy	No	No	n/d
Faerber <i>et al.</i> ^[56]	5	Cough, tachypnoea and wheezing for 2 years	CXR, chest US + CT	No	Tumour resection	n/d	No	No	n/d
Furukawa <i>et al.</i> ^[57]	3	LRTI, CXR	CT chest + MRI	No	Tumour resection	Thoracoscopy	No	No	No
Gregory <i>et al.</i> ^[58]	9	Incidental on CXR for asthma	CT chest + MRI	No	Thymectomy	Sternotomy	No	No	No
Gunnels <i>et al.</i> ^[59]	4	Incidental CXR	Cardiac catheterisation	No	n/d	Sternotomy	No	No	No
Gupta <i>et al.</i> ^[510]	11	Chest pain	CXR + CT	EUS-FNA	Tumour resection	Thoracotomy	No	No	No
Herrera <i>et al.</i> ^[511]	8	Incidental CXR	Bronchogram, aortogram, CT scan	No	Tumour resection	Thoracotomy	No	No	n/d
Kitano <i>et al.</i> ^[512]	6	Bulging thorax	CXR + CT + MRI + angiography	No	Tumour resection	Sternotomy	No	No	n/d
Alizadeh <i>et al.</i> ^[513]	6	Persistent cough	CXR + CT	EUS FNA + open	Tumour resection	Sternotomy + thoracotomy	No	No	No
Nishimura <i>et al.</i> ^[514]	6	Incidental CXR	CT chest + MRI	FNA	Tumour resection	n/d	No	No	No
Onoda <i>et al.</i> ^[515]	2	Stridor	CXR + CT + MRI	No	Tumour resection	n/d	No	No	No
Otto <i>et al.</i> ^[516]	5	Anaemia + recurrent infections (erythrocyte hypoplasia and hypogammaglobulinaemia)	CXR	n/d	n/d	n/d	n/d	No	No
Park <i>et al.</i> ^[517]	16	n/d	CT scan	n/d	Tumour resection	Robotic subxiphoid	n/d	No	No
Parakh <i>et al.</i> ^[5]	6 mo	Tachypnoea	CXR + CT scan	US-guided TruCut	Tumour resection	Thoracotomy	No	No	n/d
Patel <i>et al.</i> ^[518]	11	Right cervical mass	CXR + CT scan	Open	Tumour resection	Sternotomy	No	No	No
Ringe <i>et al.</i> ^[519]	17	Incidental CXR	CXR	No	Tumour resection	Sternotomy	No	No	No

(continued)

Table 1. (continued) Summary of a literature review of paediatric thymolipomas (references available as a supplementary file at <https://www.samedical.org/file/1873>)

Author	Age (yrs)	Presentation	Investigation	Biopsy	Treatment	Approach	Invasion	Malignancy	Complications
Rosado-de-Christenson <i>et al.</i> ^[S20]	0 - 19	n/d	n/d	n/d	n/d	n/d	n/d	n/d	n/d
Shillitoe and Goodyear ^[S21]	7	LRTI, CXR	CXR	No	Thymectomy	n/d	No	No	No
Andrus and Foot ^[S22]	13	Dyspnoea	n/d	n/d	n/d	n/d	n/d	n/d	n/d
Bigelow and Ehler ^[S23]	10	Cough	n/d	n/d	n/d	n/d	n/d	n/d	n/d
Tan <i>et al.</i> ^[S24]	4	Asthma, CXR	CT scan	No	Tumour resection	Thoracoscopy	No	No	No

CXR = chest radiograph; CT = computed tomography; FNA = fine-needle aspiration; MRI = magnetic resonance imaging; LRTI = lower respiratory tract infection; n/d = not documented in the case report; US = ultrasound; EUS = endoscopic ultrasound.



Fig. 1. (A) Frontal chest radiograph showing right homogeneous opacification in the middle and lower zones including the left middle zone. (B) Lateral view indicating anterior mediastinal mass (arrow).




Fig. 2. (A) Axial cut mediastinal window computed tomography scan of the chest showing the soft-tissue dense mass on the right, herniating to the left hemithorax (black arrow). The blue arrow shows the compressed right bronchus. (B) Coronal view showing the compressed right lung (red arrow).



Fig. 3. Anterior sternotomy revealed the well-capsulated mass, which was resected in totality.

review, a preoperative biopsy specimen was obtained in only 6/36 patients (16.2%), 4 via fine-needle aspiration, 1 open and 1 Tru-Cut. In our case, a Tru-Cut biopsy revealed that

the tissue was fatty in nature and ruled out malignant disease. We believe that obtaining a preoperative diagnosis is important in planning surgical intervention. Ruling out

malignancy allows the surgeon to avoid pursuing an aggressive, radical procedure, and even to consider minimally invasive approaches. The fact that no recurrence or malignant transformation has been reported supports limited tumour excision through a minimally invasive approach.^[6]

The definitive treatment for thymolipomas is surgical.^[1] Several surgical approaches have been used, including thoracotomy, video-assisted thoracoscopy and sternotomy, and the decision depends on the size and location of the tumour.^[1] In our review, the approach was reported in only 16 cases, with sternotomy being the most commonly used, followed by thoracotomy. The approach in our case was dictated by the size of the tumour, as a sternotomy allows good access to the mediastinum and both hemithoraces, and is usually well tolerated in children. Published data on the paediatric population reveal no malignant or surgical complications after intervention.

Conclusion

Thymolipomas are rare, slow-growing benign neoplasms, mostly diagnosed incidentally, and should be considered as a differential diagnosis when a child presents with an anterior mediastinal mass. Compression symptoms indicate a tumour of large size. Plain radiographs can be misleading in some cases, or delay definitive diagnosis and management. Surgical intervention remains the only curative measure, and is tolerated well in the paediatric population.

Declaration. None.

Acknowledgements. None.

Author contributions. Equal contributions.

Funding. None.

Conflicts of interest. None.

1. Aghajanzadeh M, Asgary MR, Mesbah A, et al. Giant thymolipoma of mediastinum and neck – initially misdiagnosed as liposarcoma by core needle biopsy. *J Fam Med Prim Care* 2018;7(5):1079-1082. https://doi.org/10.4103/jfmpc.jfmpc_228_17
2. Hall GF. A case of thymolipoma with observations on a possible relationship to intrathoracic lipomata. *Br J Surg* 1949;36(143):321-324. <https://doi.org/10.1002/bjs.18003614318>
3. Othman SA, Alfrayyan OY, Alghamdi ZM, et al. Thymolipoma association with myasthenia gravis: Case report. *Am J Case Rep* 2020;21:e923989. <https://doi.org/10.12659/ajcr.923989>
4. Moran CA, Rosado-de-Christenson M, Suster S. Thymolipoma: Clinicopathologic review of 33 cases. *Mod Pathol* 1995;8(7):741-744.
5. Parakh A, Singh V, Subramaniam R, Narula MK, Agarwala SK, Shukla S. Giant thymolipoma in an infant. *Paediatr Int Child Health* 2014;34(3):230-232. <https://doi.org/10.1179/2046905513Y.0000000108>
6. Vaziri M, Rad K. Progressive dyspnea in a 40-year-old man caused by giant mediastinal thymolipoma. *Case Rep Surg* 2016;2016:3469395. <https://doi.org/10.1155/2016/3469395>
7. Nishino M, Ashiku SK, Kocher ON, Thurer RL, Boisselle PM, Hatabu H. The thymus: A comprehensive review. *Radiographics* 2006;26(2):335-348. <https://doi.org/10.1148/rg.262045213>

Accepted 8 July 2022.