

The CPD programme for SAJCH is administered by Medical Practice Consulting.
CPD questionnaires must be completed online at www.mpconsulting.co.za

Breastfeeding intentions and behaviours of doctor mothers in Bloemfontein, South Africa

1. The Code of Good Practice on the Protection of Basic Conditions of Employment and Pregnancy in SA (1998) states that two 30-minute breaks each day for expressing should be allowed. This should continue for the child's:
 - a. First 3 months.
 - b. First 6 months.
 - c. First 9 months.
 - d. First 12 months.
2. Of the 71% of respondents in the study that did not have a dedicated room at work where they could express breastmilk, many of them experienced the following problem:
 - a. Someone had walked in on them expressing breastmilk.
 - b. Someone forbid them to express breastmilk.
 - c. Someone told them to go somewhere else to express breastmilk.
 - d. Someone encouraged them to continue to express breastmilk.

Clinical, biochemical, and histopathological diagnosis of Egyptian paediatric patients with suspected mitochondrial diseases: A hospital-based study

3. For appropriate diagnosis of mitochondrial enzymes, it is essential to follow a step-wise approach, including
 - a. measurement of the individual mitochondrial enzymes
 - b. histochemical enzymes assay in corresponding tissue
 - c. molecular diagnosis of most common known mutations
 - d. all of the above
4. The scoring criteria used for suspected patients' selection followed
 - a. Morava scoring
 - b. Walker criteria
 - c. Bernier criteria
 - d. None of the above

The value of routine empiric antibiotic use in neonates born to mothers with prolonged rupture to membranes.

5. Which one of the following statements is correct regarding proven maternal risk factors for neonatal sepsis?
 - a. Maternal fever of 37.8.
 - b. Group B Strep colonisation
 - c. Previous neonate infected with Group B Strep.
 - d. B and C

Outcome of infants with necrotising enterocolitis at Charlotte Maxeke Johannesburg Academic Hospital, South Africa

6. Which of the following variables was/is not associated with increased mortality among infants with necrotising enterocolitis in this study?
 - a. lower birth weight
 - b. HIV exposure
 - c. invasive ventilation
 - d. major birth defect

Crescentic glomerulonephritis in children: A retrospective review of data from Chris Hani Baragwanath Academic Hospital

7. Which is an example of immune-complex-mediated CGN?
 - a. Lupus nephritis
 - b. Post infectious glomerulonephritis
 - c. ANCA-associated vasculitis
 - d. Anti-GBM disease
8. What is an important bedside investigation to be done in patients who present in haematuria?
 - a. blood pressure
 - b. Serum creatinine
 - c. Renal ultrasound
 - d. Renal biopsy

Asthma control among adolescents in the inner provinces of South Africa: Perception v. reality

9. The following factors have been attributed to poor control of asthma.
 - a. Lack of disease awareness.
 - b. Medication fears.
 - c. Affordability and access to asthma therapy.
 - d. Stigma associated with taking asthma medication.
10. Regarding the Asthma Control Test, which of the following statements are true?
 - a. The ACT is a validated, self-administered.
 - b. It is 4-item questionnaire that assesses the level of asthma control.
 - c. It assesses the level of asthma control over a period of 4 weeks.
 - d. An ACT score between 15 and 25 suggests well controlled asthma.

A maximum of 3 CEUs will be awarded per correctly completed test.
CPD questionnaires must be completed online via www.mpconsulting.co.za.
After submission you can check the answers and print your certificate.
Accreditation number: MDB015/MPDP/038/027 (Clinical)